

The Impact of Charcoal Production for Energy on Tropical Rainforest Resources in Nigeria

Citation for published version (APA):

Lansu, A., Bos, J., & Ivens, W. (Accepted/In press). *The Impact of Charcoal Production for Energy on Tropical Rainforest Resources in Nigeria*. Abstract from European Geosciences Union General Assembly 2020, Vienna, Austria. <https://doi.org/10.5194/egusphere-egu2020-11780>

DOI:

[10.5194/egusphere-egu2020-11780](https://doi.org/10.5194/egusphere-egu2020-11780)

Document status and date:

Accepted/In press: 31/01/2020

Document Version:

Early version, also known as pre-print

Document license:

CC BY

Please check the document version of this publication:

- A submitted manuscript is the version of the article upon submission and before peer-review. There can be important differences between the submitted version and the official published version of record. People interested in the research are advised to contact the author for the final version of the publication, or visit the DOI to the publisher's website.
- The final author version and the galley proof are versions of the publication after peer review.
- The final published version features the final layout of the paper including the volume, issue and page numbers.

[Link to publication](#)

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal.

If the publication is distributed under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license above, please follow below link for the End User Agreement:

<https://www.ou.nl/taverne-agreement>

Take down policy

If you believe that this document breaches copyright please contact us at:

pure-support@ou.nl

providing details and we will investigate your claim.

Downloaded from <https://research.ou.nl/> on date: 30 Mar. 2023

Open Universiteit
www.ou.nl



EGU2020-11780

EGU General Assembly 2020

© Author(s) 2020. This work is distributed under the Creative Commons Attribution 4.0 License.



The Impact of Charcoal Production for Energy on Tropical Rainforest Resources in Nigeria

Angelique Lansu, Jaap Bos, and Wilfried Ivens

Open University of the Netherlands, Faculty of Science, Environmental Sciences, Heerlen, Netherlands (ala@ou.nl)

In Sub Saharan Africa, many people depend on biomass for their household energy. Charcoal production is a common technique for converting biomass into a useful energy source. Nigeria is the biggest charcoal producer in Sub Saharan Africa. A large amount of wood is harvested from Nigerian forests for this charcoal production for energy. The Nexus of charcoal-land use change-energy imposes a considerable burden on the amount of wood that must be extracted from the forest for charcoal production. Therefore, charcoal production is linked to deforestation and forest degradation. However, it is not clear to what extent the demand for charcoal in Nigeria contributes to deforestation by land use change, and degradation of forests by selected wood logging. In this study, an attempt was made to provide an answer to this and to state which situation could occur by 2030, following the expected population growth in Nigeria. To achieve this, literature and open data on charcoal production, deforestation, forest degradation and population growth in Nigeria have been collected and analysed. Subsequently, calculations were carried out to determine to what extent charcoal production contributed to deforestation in the period 1990-2015. In this period, the share of deforestation due to charcoal production increased from 6% to 14%. If the expected charcoal production in 2030 were to apply to the current situation, this share would be around 20%. The quantity of wood required can also be expressed in numbers of hectares with biomass. In that case, around 80,000 ha would be required in 2030. To validate the findings, further research is needed on the amount of biomass per hectare in Nigerian forests, and on the amount of charcoal exported, not only as source of household energy but also globally as barbecue fuel. A more extensive analysis of open data on the nexus charcoal-land use change-energy at multiple scales will help to project future interlinkages.