

# From Pasture and Cropland to Nut Orchards

## Citation for published version (APA):

Roest, E., Lansu, A., Baltissen, T., & Dekker, S. C. (2020). *From Pasture and Cropland to Nut Orchards: Modelling the Dynamics of Carbon Sequestration by Agroforestry Systems in the Temperate Climate Zone*. Abstract from European Geosciences Union General Assembly 2020, Vienna, Austria.  
<https://meetingorganizer.copernicus.org/EGU2020/EGU2020-11195.html>

## Document status and date:

Published: 01/05/2020

## Please check the document version of this publication:

- A submitted manuscript is the version of the article upon submission and before peer-review. There can be important differences between the submitted version and the official published version of record. People interested in the research are advised to contact the author for the final version of the publication, or visit the DOI to the publisher's website.
- The final author version and the galley proof are versions of the publication after peer review.
- The final published version features the final layout of the paper including the volume, issue and page numbers.

[Link to publication](#)

## General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal.

If the publication is distributed under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license above, please follow below link for the End User Agreement:

<https://www.ou.nl/taverne-agreement>

## Take down policy

If you believe that this document breaches copyright please contact us at:

[pure-support@ou.nl](mailto:pure-support@ou.nl)

providing details and we will investigate your claim.

Downloaded from <https://research.ou.nl/> on date: 16 May. 2022

Open Universiteit  
[www.ou.nl](http://www.ou.nl)



# From Pasture and Cropland to Nut Orchards

## Citation for published version (APA):

Roest, E., Lansu, A., Baltissen, T., & Dekker, S. C. (Accepted/In press). *From Pasture and Cropland to Nut Orchards: Modelling the Dynamics of Carbon Sequestration by Agroforestry Systems in the Temperate Climate Zone*. Abstract from European Geosciences Union General Assembly 2020, Vienna, Austria.

## Document status and date:

Accepted/In press: 31/01/2020

## Please check the document version of this publication:

- A submitted manuscript is the version of the article upon submission and before peer-review. There can be important differences between the submitted version and the official published version of record. People interested in the research are advised to contact the author for the final version of the publication, or visit the DOI to the publisher's website.
- The final author version and the galley proof are versions of the publication after peer review.
- The final published version features the final layout of the paper including the volume, issue and page numbers.

[Link to publication](#)

## General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal.

If the publication is distributed under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license above, please follow below link for the End User Agreement:

<https://www.ou.nl/taverne-agreement>

## Take down policy

If you believe that this document breaches copyright please contact us at:

[pure-support@ou.nl](mailto:pure-support@ou.nl)

providing details and we will investigate your claim.


Downloaded from <https://research.ou.nl/> on date: 06 Jan. 2021

Open Universiteit  
[www.ou.nl](http://www.ou.nl)



Session SSS9.7 (D2112)

# *From Grassland and Cropland to Nut Orchards: Carbon Sequestration Dynamics of temperate Agroforestry Systems*



**EGU2020-11195**

**presentation**

6 mei 2020 (8:30-10:15)

Erik Roest<sup>1</sup>, co-authors: Angelique Lansu<sup>1</sup>, Ton Baltissen<sup>13</sup> and Stefan Dekker<sup>12</sup>

<sup>1</sup>Open University Heerlen (NL), <sup>2</sup>Utrecht University, <sup>3</sup>Cropeye

**Open Universiteit**

[www.ou.nl](http://www.ou.nl)



# Introduction



The general discourse is that agroforestry systems (AFS) can sequester more carbon than crop- or grassland (Pardon et al., 2017).

Study	Country (AEZ)	Land use	C-sequestration rate (Mg C·ha <sup>-1</sup> yr <sup>-1</sup> )	Range
Cardinael et al. (2017)	France (temperate)	silvoarable to regular AM	0.69	vegetation and soil
Hamon et al. (2009) as cited in Aertsens et al. (2013)	Europe (mainly temperate)	regular AM to agroforestry (e.g. based on Juglans)	1.5-4.0 [2.75]	vegetation and soil
Pardon et al. (2017)	Belgium (temperate)	regular AM to silvoarable	0.21	soil
Sharrow and Ismail (2004)	Oregon, USA (temperate)	pasture to silvopastoral	0.52	vegetation and soil
Wotherspoon et al. (2014)	Ontario, Canada (temperate)	regular AM to silvoarable (various tree species)	0.8–2.1	vegetation and soil

## Research question:

To what extent does changing crop- or grassland into nut orchards in the temperate zone contribute to increased carbon sequestration in vegetation and soil?


Open Universiteit  
[www.ou.nl](http://www.ou.nl)




(Roest et al., 2020)

# Method (1)

- Object of study: orchards of *Corylus* (hazelnut trees) and solitary trees of *Juglans* (walnut trees) in the province of Gelderland, The Netherlands.
- The research was conducted with the use of chronosequences.
- The basis of our study was to depict an overview of C-stocks & –fluxes in nut orchards.



# Method (2)




Carbon stocks in terrestrial ecosystems (partly based on Lal (2005)).

Our research concentrated on three main stocks of carbon: soil organic carbon (SOC), belowground biomass (BGB) and aboveground biomass (AGB) and the fluxes that run from one to another.


# Results (1)

- C-stocks in *Corylus* and *Juglans* orchards are larger than C-stocks in reference parcels (used as grassland or cropland).
- The figure below shows the C-stocks in the four chronosequences we distinguished, with on the left the reference parcel and on the right the parcel with the oldest trees.



# Results (2)

- C-stocks in orchards show a positive correlation with tree age.
- In general carbon in soil (SOC) and in biomass (aboveground biomass and belowground biomass) have an almost similar contribution to the additional C- sequestration (the figure below shows the SOC-stock in the four different sequences).
- Our results on C-fluxes are largely in line with sequestration rates found in various studies on agroforestry systems in the temperate zone.





# Discussion & conclusions



- Changing crop- or grassland into *Corylus* and *Juglans* orchards in the temperate zone can attribute to increased C-sequestration (wood harvest included).

C-flux	Unit	<i>Corylus</i>	<i>Juglans</i>
SOC	Mg C ha <sup>-1</sup> y <sup>-1</sup>	-/-0.12-2.16	0.35-1.48
BGB	Mg C ha <sup>-1</sup> y <sup>-1</sup>	0.04-0.41	0.02-0.14
AGB	Mg C ha <sup>-1</sup> y <sup>-1</sup>	0.53-1.15	0.25-0.67
Total	Mg C ha <sup>-1</sup> y <sup>-1</sup>	0.82-3.36	1.16-1.75

- Composing uniform chronosequences is difficult.
- Changing crop- or grassland into *Corylus* and *Juglans* orchards seems promising to mitigate part of the anthropogenic CO<sub>2</sub> emissions to the atmosphere.
- Additional research on sequestration rates, preferably by time studies, is recommended.



# References and colofon



## References

- Cardinael, R., Chevallier, T., Cambou, A., Béral, C., Barthès, B. G., Dupraz, C., . . . Chenu, C. (2017). Increased soil organic carbon stocks under agroforestry: A survey of six different sites in France. *Agriculture, Ecosystems and Environment*, 236, 243-255. doi:10.1016/j.agee.2016.12.011
- Aertsens, J., De Nocker, L., & Gobin, A. (2013). Valuing the carbon sequestration potential for European agriculture. *Land Use Policy*, 31, 584-594. doi:10.1016/j.landusepol.2012.09.003
- Pardon, P., Reubens, B., Reheul, D., Mertens, J., De Frenne, P., Coussement, T., . . . Verheyen, K. (2017). Trees increase soil organic carbon and nutrient availability in temperate agroforestry systems. *AGRICULTURE ECOSYSTEMS & ENVIRONMENT*, 247, 98-111. doi:10.1016/j.agee.2017.06.018
- Sharrow, S. H., & Ismail, S. (2004). Carbon and nitrogen storage in agroforests, tree plantations, and pastures in western Oregon, USA. *Agroforestry Systems*, 60(2), 123-130. doi:10.1023/B:AGFO.0000013267.87896.41
- Wotherspoon, A., Thevathasan, N. V., Gordon, A. M., & Voroney, R. P. (2014). Carbon sequestration potential of five tree species in a 25-year-old temperate tree-based intercropping system in southern Ontario, Canada. *Agroforestry Systems*, 88(4), 631-643. doi:10.1007/s10457-014-9719-0
- Lal, R. (2005). Forest soils and carbon sequestration. *Forest Ecology and Management*, 220(1), 242-258. doi:10.1016/j.foreco.2005.08.015

## Colofon

Financing of all laboratory analysis came from the Province of Gelderland, The Netherlands under the project named 'Notenbomen als toplaagverbeteraar'. The project 'Notenbomen als toplaagverbeteraar' was initiated and led by Ton Baltissen from the company Cropeye.



Open Universiteit  
www.ou.nl



(Roest et al., 2020)