

The relationship between the global mean deep-sea and surface temperature during the Early Eocene

Citation for published version (APA):

Goudsmit, B., Lansu, A., von der Heydt, A. S., Zhang, Y., & Ziegler, M. (2022). *The relationship between the global mean deep-sea and surface temperature during the Early Eocene*. 1. Abstract from European Geosciences Union General Assembly 2022, Vienna, Austria. <https://doi.org/10.5194/egusphere-egu22-9897>

DOI:

[10.5194/egusphere-egu22-9897](https://doi.org/10.5194/egusphere-egu22-9897)

Document status and date:

Published: 01/05/2022

Document Version:

Peer reviewed version

Document license:

CC BY

Please check the document version of this publication:

- A submitted manuscript is the version of the article upon submission and before peer-review. There can be important differences between the submitted version and the official published version of record. People interested in the research are advised to contact the author for the final version of the publication, or visit the DOI to the publisher's website.
- The final author version and the galley proof are versions of the publication after peer review.
- The final published version features the final layout of the paper including the volume, issue and page numbers.

[Link to publication](#)

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal.

If the publication is distributed under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license above, please follow below link for the End User Agreement:

<https://www.ou.nl/taverne-agreement>

Take down policy

If you believe that this document breaches copyright please contact us at:

pure-support@ou.nl

providing details and we will investigate your claim.

Downloaded from <https://research.ou.nl/> on date: 21 Jun. 2024

Open Universiteit
www.ou.nl



EGU22-9897

EGU General Assembly 2022

© Author(s) 2022. This work is distributed under the Creative Commons Attribution 4.0 License.



The relationship between the global mean deep-sea and surface temperature during the Early Eocene

Barbara Goudsmit^{1,2}, Angeliqne Lansu³, Anna S. von der Heydt⁴, and Martin Ziegler²

¹Department of Estuarine & Delta Systems, NIOZ Royal Netherlands Institute for Sea Research, Yerseke, The Netherlands

²Department of Earth Sciences, Utrecht University, Utrecht, The Netherlands

³Department of Environmental Sciences, Open Universiteit, Heerlen, Netherlands

⁴Institute for Marine and Atmospheric Research, Utrecht University, Utrecht, The Netherlands

Under continued high anthropogenic CO₂ emissions, the atmospheric CO₂ concentration around 2100 will be like that of the Early Eocene Climate Optimum (EECO, 56–48 Ma) hothouse period. Hence, reconstructions of the EECO climate give insight into the workings of the climate system under the possible future CO₂ conditions. Our current understanding of global mean surface temperature (GMST) during the Cenozoic era relies on paleo-proxy estimates of deep-sea temperature (DST) combined with assumed relationships between global mean DST (GMDST), global mean sea-surface temperature (GMSST), and GMST. The validity of these assumptions is essential in our understanding of past and future climate states under hothouse conditions.

We analyse the relationship between these global temperature indicators for the end-of-simulation global mean temperature values in 25 different millennia-long model simulations of the EECO climate under varying CO₂ levels, performed as part of the Deep-Time Model Intercomparison Project (DeepMIP). The model simulations show limited spatial variability in DST, indicating that local DST estimates can be regarded representative of GMDST. Linear regression analysis indicates that GMDST and GMST respond stronger to changes in atmospheric CO₂ than GMSST by factors 1.18 and 1.17, respectively. Consequently, the responses of GMDST and GMST to atmospheric CO₂ changes are similar in magnitude. This model-based analysis indicates that changes in GMDST can be used to estimate changes in GMST during the EECO, validating the assumed relationships. To test the robustness of these results, other Cenozoic climate states besides EECO should be analysed similarly.