

Is extreme sexually behaviour a specific behavioural disorder of dementia or not?

Citation for published version (APA):

Waterink, W., Bartelet, M., & Van Hooren, S. (2013). *Is extreme sexually behaviour a specific behavioural disorder of dementia or not?*. Poster session presented at The 8th International Congress on Vascular Dementia & The First Cognitive Impairment European Meeting, Athens, Greece.

Document status and date:

Published: 01/01/2013

Document Version:

Publisher's PDF, also known as Version of record

Please check the document version of this publication:

- A submitted manuscript is the version of the article upon submission and before peer-review. There can be important differences between the submitted version and the official published version of record. People interested in the research are advised to contact the author for the final version of the publication, or visit the DOI to the publisher's website.
- The final author version and the galley proof are versions of the publication after peer review.
- The final published version features the final layout of the paper including the volume, issue and page numbers.

[Link to publication](#)

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal.

If the publication is distributed under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license above, please follow below link for the End User Agreement:

<https://www.ou.nl/taverne-agreement>

Take down policy

If you believe that this document breaches copyright please contact us at:

pure-support@ou.nl

providing details and we will investigate your claim.

Downloaded from <https://research.ou.nl/> on date: 09 Aug. 2022

Open Universiteit
www.ou.nl



IS EXTREME SEXUALLY BEHAVIOUR A SPECIFIC BEHAVIOURAL DISORDER OF DEMENTIA OR NOT?

Wim Waterink¹, Marjukka Bartelet¹, & Susan van Hooren^{1,2}

¹Open University of the Netherlands, Heerlen; ²Zuyd University of Applied Sciences, Heerlen

Introduction

In nursing homes, extreme sexually behaviour is one of the most challenging behaviours in dementia. However, in the literature there is no conformity in how to label and define this type of behaviour. According to recent theoretical perspectives, extreme sexually behaviour may be regarded as a part of disinhibited behaviour or could be considered as an independent neuropsychiatric symptom. In this multicentre study, it was investigated whether there is a relationship between extreme sexually behaviour and typical neuropsychiatric symptoms seen in dementia.

Methods and Materials

In 179 residents diagnosed with dementia, extreme sexually behaviour was measured using an observation scale. Twelve neuropsychiatric symptoms were measured by the Neuropsychiatric Inventory (NPI).

Table 1.
Comparisons between residents with (n=43) and without (n=136) observed extreme sexually behaviour (ESB) on the 12 NPI disorders using MANOVA

	With ESB		Without ESB		p	η^2
	M (SD)	M (SD)	F(1,176)	p		
Delusions	1.91 (3.76)	1.53 (2.92)	.004	.951	.000	
Hallucinations	1.16 (3.22)	.88 (2.67)	.126	.723	.001	
Agitation/Aggression	4.12 (4.17)	3.33 (4.08)	.091	.763	.001	
Depression/Dysphoria	2.70 (3.99)	1.32 (2.80)	3.295	.071	.018	
Anxiety	1.72 (3.24)	1.57 (2.81)	.250	.618	.001	
Elation/Euphoria	.74 (1.99)	.25 (1.05)	2.856	.093	.016	
Apathy/Indifference	2.67 (4.08)	2.68 (4.26)	.418	.519	.002	
Disinhibition	3.88 (3.86)	.88 (2.41)	27.279	.000	.134	
Irritability/Lability	2.86 (3.49)	2.81 (3.76)	.455	.501	.003	
Motor disturbance	2.35 (3.79)	2.26 (3.65)	.000	.987	.000	
Sleep disturbance	2.16 (3.77)	1.13 (2.59)	.577	.449	.003	
Eating disturbance	1.19 (2.62)	1.88 (3.34)	3.370	.068	.019	

Results

Multivariate Analysis of Covariance with gender, showed that residents with observed extreme sexually behaviour only showed a higher score on neuropsychiatric symptom 'disinhibition', as compared to residents with non-observed sexually behaviour. In addition, the effect size was large.

However, this difference could be biased by the fact that the subscale had two specific items on sexual behaviour. Therefore, a corrected NPI-disinhibition variable was created, by removing these items from the original subscale. A follow-up regression analyses was performed, with gender as a covariate.

Table 2
Extreme sexually behaviour (ESB) a predictor of corrected NPI-disinhibition

	corrected NPI-disinhibition	
	ΔR^2	B
Step 1	.068***	
Constant		2.214***
Gender		-.822***
Step 2	.063***	
Constant		1.474**
Gender		-.474
ESB		.138***
Total R ²	.132***	
n	179	

Note. p < 0,01. *** p < 0,001.

Conclusion

These findings indicate that among residents with dementia, extreme sexually behaviours should not be considered as an independent neuropsychiatric symptom. Instead, disinhibition may be an important underlying mechanism for extreme sexually behaviour and thus validates the label 'sexually disinhibited behaviour'.

